

Male genital eczema factsheet

With any genital condition it is important that you have the correct diagnosis made by your doctor so that any possible underlying conditions can be identified and treated. Please do not feel embarrassed about asking for a proper examination to be performed.

There are different types of eczema that could affect the genital area, including atopic eczema, seborrhoeic dermatitis, allergic contact eczema and irritant contact eczema (for more information on the different types of eczema, please visit the NES website or contact our Helpline).

Genital eczema frequently affects the penis, scrotum, groin, the skin around the opening of the anus and the skin between the cheeks of the buttocks (perianal eczema). There are other causes of red, sore skin, such as fungal and yeast infections and sexually transmitted infections (STIs), which might need to be ruled out by your doctor. Your GP might also refer you to dermatology, particularly if allergic contact dermatitis is suspected or diagnosis is uncertain.

If you have genital eczema, it is not always possible to discover the cause, although sometimes it can be due to a contact allergy. However, genital skin can be particularly prone to irritation because of sweating, tight clothing, and friction between skin surfaces. Soaps, bubble baths, shower gels, talcum powder, moist toilet tissue/wipes, aftershave, personal deodorants, haemorrhoid preparations, antiseptics such as Dettol, Savlon and TCP, and similar products can also trigger eczema. Urine and faeces are especially irritating to the skin, and incontinence always makes any genital condition worse and harder to treat. If you are incontinent of urine, faeces, or both, discuss this with your doctor or nurse, who can refer you to specialist teams. Trying to manage the problem with wipes or pads may make the eczema worse. Don't feel embarrassed about it.

Treatment

If you experience genital irritation, it is better to seek medical help than to self-medicate, since some over-the-counter treatments contain potentially allergenic and

irritant ingredients. Your doctor will usually prescribe emollients and topical steroids to treat genital and perianal eczema.

Emollients can be applied to the genital area as required. They should be reapplied after bathing and showering. Use emollient as a soap substitute and avoid all soap and cosmetic washes. It is also a good idea to wash with emollients after opening your bowels to prevent infection when skin is red and sore.

Steroids are safe to use in the genital area as long as they are of the correct strength and are used appropriately. The skin here can absorb topical steroids more readily than in other parts of the body, and topical steroids should therefore be used carefully in the genital area. They only need to be used once or twice a day, as prescribed, and a 30g tube should normally last at least 3 months. It is important to avoid prolonged or over-use of combination steroid preparations, in particular those containing certain antibiotics such as neomycin, which may cause contact dermatitis. If combined topical

Male genital eczema factsheet

steroids and antibiotic creams are prescribed for infection, they should be used for a maximum of 14 days, after which you should return to plain topical steroids if the eczema is still flaring.

Ointment-based topical treatments contain fewer potential allergens than creams, so are especially suitable for sensitive areas. Since ointments are greasy, they generally spread easily and are well-absorbed. However, creams are easier to spread on hair-bearing skin.

If using a topical steroid, apply this at least half an hour before using an emollient (please see the NES factsheets on Emollients and Topical Steroids).

Itching

Genital eczema can cause embarrassment because the area is intensely itchy, leading to scratching. Anti-itch preparations, including anti-itch emollients (Balneum Plus or E45 Itch) may be helpful in relieving severe itching and discomfort. Ask the pharmacist if you are uncertain about a product's ingredients. You can keep your emollients in the fridge, as a cold application can be more soothing.

Some people develop an allergy to nail varnish, so if your partner wears it and you think this may be causing problems for you, see if leaving it off makes a difference.

If the itching persists, don't try to treat yourself – seek medical advice. You may be offered a blood test if, by reference to your medical history, your healthcare professional thinks it necessary – for example, ferritin (iron) levels might be checked as iron deficiency anaemia can cause itching; or you may need further tests such as patch testing to exclude allergic contact dermatitis.

If night-time itching is a problem, an antihistamine may help you to sleep.

Remember, if you have genital eczema, itching may also occur due to yeast infections (thrush) or STIs, which can cause intense itch and invariably make eczema worse. If you have any different symptoms to eczema, do ask a health care professional for further advice.

Hygiene

It is important to keep the genital area clean. However, washing the area too much may encourage further irritation, especially if abrasive sponges or flannels are used. Uncircumcised males should pull back the foreskin in order to wash the head of the penis.

Wash with warm water and an emollient soap substitute or use your usual leave-on emollient cream for washing. Avoid soaps, antiseptic washes, moist tissue wipes, bubble bath, deodorant, perfume, fragrance and any other products that could irritate and dry out the skin. Also, avoid using shampoo in the bath or shower since it could inadvertently come into contact with the genital area and cause irritation. If you do need to shampoo your hair in the bath, apply your emollient cream to the genital skin first and then rinse it off after you have rinsed your hair.

Good hand hygiene is important when using emollients and topical steroids – always wash your hands before and after applying treatments. If possible, use emollients from a pump dispenser. If you are using cream or ointment from a pot, do not put your hands in and out of the pot as there is a risk of infection – instead, use a clean spoon or spatula to decant the amount you need to use from the pot.

If you wish to remove hair in the genital area, waxing and shaving products may irritate the skin further. Using your emollient wash product may be a more gentle option on the skin (and while you're there, it's a good idea to check your testicles for lumps once a month after a warm bath or shower).

Clothing

Comfortable, loose cotton clothing will help to lessen irritation and friction. Wear loose-fitting underwear (100% cotton or silk is more comfortable, as natural fibres will not be irritating). Avoid thongs. Most people prefer to use a non-biological washing powder to wash their clothes. Finding the most suitable one is a case of trial and error. You may find that you need to wash at 60°C or higher with the non-biological powder to clean your clothing effectively and remove the grease from emollients. Always ensure that your clothing is rinsed well. It is best to avoid fabric conditioners since they contain fragrance and other ingredients that might cause irritation or an allergy.

Piercing and tattoos

Body piercing does not cause eczema, but problems can occur later on as a consequence. Nickel allergy is very common and the chance of this developing is greatly increased if you have body piercing and use cheap metal jewellery. Having a tattoo (particularly when black henna or dye is used) increases the risk of developing a contact allergic dermatitis to dyes especially PPD (p-Phenylenediamine). There is also a high risk of pierced and tattooed areas becoming infected.

Sex

Be careful not to use topical steroids shortly before having sex on an area that is likely to transfer them onto your partner. You need to let the topical steroids be completely absorbed into the genital area first. As long as you do this, topical steroids should not cause any problems. The same applies to oral sex.

Sperm fluid (semen) can make irritant eczema worse; however, a true allergic reaction to sperm or sperm fluid, which in its severest form can cause an anaphylactic reaction, is exceptionally rare.

Barrier contraceptives

Most condoms, diaphragms and caps are made from latex rubber and generally do not cause problems for people with eczema. However, some people have an allergy to latex or other rubber ingredients and they should use products made from material such as polyurethane or silicone.

Spermicides can also irritate the skin. If this is a problem or concern, discuss this with your pharmacist, doctor, or staff at your family planning clinic.

Emollients and topical steroid creams can affect the integrity of condoms and diaphragms, so be aware that their effectiveness will be reduced if creams have been applied recently.

Lubricants

Personal lubricants (or lubes) are specialized lubricants that help to reduce friction, pain and discomfort during sexual acts. There are many types available and choice is a very individual thing. Water-based lubricants are water-soluble and are the type most commonly used. They do tend to dry out during use, so you may need to reapply them or add water to reactivate them. Silicone-based lubricants are usually formulated with fewer than four ingredients and do not contain any water. Silicone lubricants feel different to water-based ones and, since they are not absorbed by skin or mucus membranes, last longer.

Conclusion

Genital eczema can be treated and controlled, but as with all types of eczema it is often a long-term condition with no cure. It is important that you use the products prescribed by your doctor or dermatologist, and attend appointments in order for the condition to be monitored and treatments changed if necessary.

Male genital eczema factsheet

Resources

National Eczema Society

www.eczema.org/factsheets

British Association of Dermatologists

www.bad.org.uk

New Zealand Dermatological Society

www.dermnetnz.org

Shah. Common skin conditions affecting the genitals

www.radcliffehealth.com/sites/radcliffehealth.com/files/books/samplechapter/0401/Shah_chpt11-442d4f80rdz.pdf

DermNet NZ (2014) Balanitis

www.dermnetnz.org/site-age-specific/balanitis.html

DermNet NZ (2013) Seborrhoeic dermatitis

www.dermnet.org.nz/dermatitis/seborrhoeic-dermatitis.html

DermNet NZ (2013) Allergic contact dermatitis

www.dermnet.org.nz/dermatitis/contact-allergy.html

Useful organisations

Brook Advisory Service

(Clinics throughout the UK for young people under 25 for contraception, sexual and relationship problems)

www.brook.org.uk

Family Planning Association

www.fpa.org.uk

Latex Allergy Support Group

www.lasg.org.uk

NHS Choices Contraception

www.nhs.uk/Conditions/contraception-guide/Pages/contraception.aspx

DISCLAIMER

Our publications contain information and general advice about eczema. They are written and reviewed by dermatology experts, with input from people with eczema. We hope you find the information helpful, although it should not be relied upon as a substitute for personalised advice from a qualified healthcare professional. While we strive to ensure the information is accurate and up-to-date, National Eczema Society does not accept any liability arising from its use. We welcome reader feedback on our publications, please email us at info@eczema.org

Factsheet last reviewed February 2018.

© National Eczema Society, June 2019. All rights reserved. Except for personal use, no part of this work may be distributed, reproduced, downloaded, transmitted or stored in any form without the written permission of National Eczema Society.



National Eczema Society is the UK charity for everyone affected by eczema. We help support people with eczema, providing information and advice, which we deliver through our website, social media, campaigns, publications and nurse-supported Helpline. We also provide a voice for people with eczema, raising awareness of the condition, supporting research into new treatments and campaigning for better medical care.

National Eczema Society is a registered charity in England and Wales (No. 1009671) and in Scotland (No. SC043669). Registered Office: 11 Murray Street, London NW1 9RE

National Eczema Society Helpline: 0800 448 0818 Email: helpline@eczema.org www.eczema.org